

For faster refills, phone in 24 hours in advance

Keep Out of Reach of Children

CVS item# 715078

CVS/pharmacy

3013 MARINA BAY DR
LEAGUE CITY, TX
77573

#7094

Ph. 281.538-2416

www.cvs.com

ARMATTA, BETH

313 WEST DR
KEMAH, TX 775650000
Ph: 832-703-7215

05/28/2011

Prscr: RAJINDER KUMAR BHALLA
Refills: 0

FUROSEMIDE 80 MG TABLET

MYLAN
TAKE 1 TABLET BY MOUTH TWICE A DAY

This is a WHITE, ROUND-shaped TABLET imprinted with MYLAN 232 on the front and 80 on the back.

FUROSEMIDE - ORAL - (fyou-ROW-seh-mide)

COMMON BRAND NAME(S):

Delone, Detue, Lasix

WARNING:

Furosemide is a very potent medication. Using too much of this drug can lead to serious water and salt/mineral loss. Therefore, it is important that you are closely monitored by your doctor while taking this medication. Tell your doctor right away if you become very thirsty or confused, or develop muscle cramps/weakness. See also Side Effects section.

USES:

Furosemide is used to reduce extra fluid in the body (edema) caused by conditions such as heart failure, liver disease, and kidney disease. This can lessen symptoms such as shortness of breath and swelling in your arms, legs, and abdomen. This drug is also used to treat high blood pressure. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Furosemide is a "water pill" (diuretic) that causes you to make more urine. This helps your body get rid of extra water and salt.

HOW TO USE:

Read the Patient Information Leaflet if available from your pharmacist before you start taking furosemide and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth as directed by your doctor, with or without food, usually once or twice daily. It is best to avoid taking this medication within 4 hours of your bedtime to prevent having to get up to urinate. Dosage is based on your medical condition, age, and response to treatment. For children, the dose is also based on weight. Older adults usually start with a lower dose to decrease the risk of side effects. Do not increase your dose or take it more often than directed. Take this medication regularly in order to get the most benefit from it. To help you remember, take it at the same time(s) of the day as directed. It is important to continue taking this medication even if you feel well. Most people with high blood pressure do not feel sick. Sucralfate, cholestyramine, and colestipol can decrease the absorption of furosemide. If you are taking any of these drugs, separate the timing of each dose from furosemide by at least 2 hours. Tell your doctor if your condition does not improve or if it worsens (for example, your blood pressure readings remain high or increase).

SIDE EFFECTS:

Dizziness, lightheadedness, headache, or blurred vision may occur as your body adjusts to the medication. If any of these effects persist or worsen, tell your doctor or pharmacist promptly. To reduce the risk of dizziness and lightheadedness, get up slowly when rising from a sitting or lying position. Remember that your doctor has prescribed this medication because he or she has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. This medication may cause a serious loss of body water (dehydration) and salt/minerals. Tell your doctor right away if you have any of these unlikely but serious side effects: muscle cramps, weakness, unusual tiredness, confusion, severe dizziness, fainting, drowsiness, unusual dry mouth/thirst, nausea, vomiting, fast/irregular heartbeat, unusual decrease in the amount of urine. Tell your doctor right away if any of these unlikely but serious side effects occur: numbness/tingling/pain/redness/swelling of the arms/legs, hearing changes (such as ringing in the ears, temporary or permanent decreased

Continued on reverse side.

IMPORTANT DISCLAIMER: The side effects listed above are not all of the possible risks that could be caused by this medication. For further information, please consult with your physician about the uses, precautions and risks of the medication.

AR
05/28/2011

PROMISED: 02:00p
05-29-2011
Scripts: 01

CVS/pharmacy #7094 Ph. 281.538-2416

3013 MARINA BAY DR
LEAGUE CITY, TX
77573

ARMATTA, BETH

313 WEST DR, KEMAH, TX 775650000
Ph. 832-703-7215 068 10 26
FUROSEMIDE 80 MG TABLET
MYLAN

TAKE 1 TABLET BY MOUTH TWICE A DAY

NDC 00378-0232-05 Days Supply: 30 Refills: 0 Qty: 60 EA

Prscr: RAJINDER KUMAR BHALLA
GR. DISNEYRX
TP. 4000
AUTHY Y11W3ET

PAY: \$4.00

PAID PRESCRIPTION

Caps: Y

002103



**Personal
Emergency
Response
System**

**FREE Equipment FREE Delivery EASY Setup
Minimal Monthly Service Fee**

Call 1-800-283-2300 for a FREE BROCHURE

www.MedicalAlertSystem.com

001489

**Your CVS Pharmacist can
answer questions about
your medications**

- Knowledgeable
- Committed
- Caring
- Confidential

CVS/pharmacy

PATIENT PRESCRIPTION INFORMATION

IF YOU HAVE ANY QUESTIONS ABOUT YOUR MEDICATION,
PLEASE CONTACT YOUR PHARMACIST:

Thu-Trang T Nghiem, RPh.